

Health Status

Life Expectancy

| Table # in OA 2000 | Table titles in OA2000 | Data source | Summary of changes between OA2000 and OA2004 | Table # in OA 2004 | Table titles in OA2004 | Data source | Summary of changes between OA2004 and OA2006 | Table # in OA 2006 | Table titles in OA2006 | Data source | Updated since OA2006 |
|--------------------|---|------------------|--|--------------------|---|------------------|--|--------------------|---|------------------|----------------------|
| 12A | Life expectancy by age group and sex, in years, 1900-97 | Vital Statistics | Updated with 2000 and 2001 data. | 13A | Life expectancy, by age and sex, selected years 1900-2001 | Vital Statistics | Updated with 2002 and 2003 data. | 13A | Life expectancy, by age and sex, selected years 1900-2003 | Vital Statistics | No |
| 12B | Life expectancy by age group and race, in years, 1997 | Vital Statistics | Updated with 2001 data and expanded. | 13B | Life expectancy, by age and race, 2001 | Vital Statistics | Updated with 2003 data. | 13B | Life expectancy, by age and race, 2003 | Vital Statistics | No |

Mortality

| | | | | | | | | | | | |
|-----|---|------------------|--|-----|---|------------------|---|-----|---|------------------|----|
| 13A | Death rates for selected leading causes of death among persons age 65 or older, 1980-97 (per 100,000) | Vital Statistics | Updated with 1998-2001 data. See note under table for change in coding information. | 14A | Death rates for selected leading causes of death among people age 65 and over, 1981-2001 | Vital Statistics | Updated with 2002 and 2003 data. Alzheimer's disease added as one of the leading causes of death. | 14A | Death rates for selected leading causes of death among people age 65 and over, 1981-2003 | Vital Statistics | No |
| 13B | Leading causes of death among persons age 65 or older, by sex and race and Hispanic origin, 1997 | Vital Statistics | Updated with 2001 data. Does not include number of deaths from all causes. | 14B | Leading causes of death among people age 65 and over, by sex and race and Hispanic origin, 2001 | Vital Statistics | Updated with 2002 data. | 14B | Leading causes of death among people age 65 and over, by sex and race and Hispanic origin, 2002 | Vital Statistics | No |
| 13C | Leading causes of death among persons age 85 or older, by sex and race and Hispanic origin, 1997 | Vital Statistics | Updated with 2001 data. Does not include number of deaths from all causes. | 14C | Leading causes of death among people age 85 and over, by sex and race and Hispanic origin, 2001 | Vital Statistics | Updated with 2002 data. | 14C | Leading causes of death among people age 85 and over, by sex and race and Hispanic origin, 2002 | Vital Statistics | No |

Chronic Health Conditions

| Table # in OA 2000 | Table titles in OA2000 | Data source | Summary of changes between OA2000 and OA2004 | Table # in OA 2004 | Table titles in OA2004 | Data source | Summary of changes between OA2004 and OA2006 | Table # in OA 2006 | Table titles in OA2006 | Data source | Updated since OA2006 |
|--------------------|---|---|--|--------------------|---|----------------------------------|---|--------------------|---|----------------------------------|----------------------|
| 14 | Percentage of persons age 70 or older who reported having selected chronic conditions, by selected characteristics, 1984 and 1995 | Supplement on Aging, Second Supplement on Aging | Changed data source from SOA I&I. Separated into two tables. Added emphysema, asthma, and chronic bronchitis. No longer have age-breaks. | 15A | Percentage of people age 65 and over who reported having selected chronic conditions, by sex, 2001-02 | National Health Interview Survey | Updated with 2003-04 data. The question used to estimate percentage of people with arthritis changed. | 15A | Percentage of people age 65 and over who reported having selected chronic conditions, by sex, 2003-04 | National Health Interview Survey | No |
| | | | New table. Changed data source from SOA I&I. Included data for five 2-year averaged periods. | 15B | Percentage of people age 65 and over who reported having selected chronic conditions, 1997-2002 | National Health Interview Survey | Updated with 2003-04 data. Note: The question used to estimate percentage of people with arthritis changed. Estimates in OA2004 are age-adjusted. Estimates in OA2006 are unadjusted. | 15B | Percentage of people age 65 and over who reported having selected chronic conditions, 1997-2004 | National Health Interview Survey | No |

Memory Impairment

| | | | | | | | | | | | |
|----|---|-----------------------------|---|----|--|-----------------------------|--------------|----|--|-----------------------------|----|
| 15 | Percentage of persons age 65 or older with moderate or severe memory impairment, by age group and sex, 1998 | Health and Retirement Study | Updated with 2002 data. Because of some changes in methods from OA2000, no inference should be made about longitudinal trends between OA2000 and OA2004. Data are only shown for the combined moderate or severe memory impairment category. | 17 | Percentage of people age 65 and over with moderate or severe memory impairment, by age group and sex, 2002 | Health and Retirement Study | Not updated. | 17 | Percentage of people age 65 and over with moderate or severe memory impairment, by age group and sex, 2002 | Health and Retirement Study | No |
|----|---|-----------------------------|---|----|--|-----------------------------|--------------|----|--|-----------------------------|----|

Depressive Symptoms

| Table # in OA 2000 | Table titles in OA2000 | Data source | Summary of changes between OA2000 and OA2004 | Table # in OA 2004 | Table titles in OA2004 | Data source | Summary of changes between OA2004 and OA2006 | Table # in OA 2006 | Table titles in OA2006 | Data source | Updated since OA2006 |
|--------------------|---|-----------------------------|--|--------------------|---|-----------------------------|--|--------------------|---|-----------------------------|----------------------|
| 16 | Percentage of persons age 65 or older with severe depressive symptoms, by age group and sex, 1998 | Health and Retirement Study | Updated with 2002 data. Different title. | 18 | Percentage of people age 65 and over with clinically relevant depressive symptoms, by age group and sex, 2002 | Health and Retirement Study | Not updated. | 18 | Percentage of people age 65 and over with clinically relevant depressive symptoms, by age group and sex, 2002 | Health and Retirement Study | No |

Self-Rated Health Status (OA2000) changed to Respondent-Assessed Health Status (OA2004 and OA2006)

| | | | | | | | | | | | |
|----|---|----------------------------------|---|----|--|----------------------------------|----------------------------|----|--|----------------------------------|----|
| 17 | Percentage of persons age 65 or older who reported good to excellent health, by age group, sex, and race and Hispanic origin, 1994-96 | National Health Interview Survey | Updated with 2000-02 data and expanded. Title changed. Different race categories. | 20 | Respondent-assessed health status among people age 65 and over, by selected characteristics, 2000-02 | National Health Interview Survey | Updated with 2002-04 data. | 20 | Respondent-assessed health status among people age 65 and over, by selected characteristics, 2002-04 | National Health Interview Survey | No |
|----|---|----------------------------------|---|----|--|----------------------------------|----------------------------|----|--|----------------------------------|----|

Disability

| | | | | | | | | | | | |
|-----|---|---|---|-----|---|-------------------------------------|---|-----|---|-------------------------------------|------|
| 18A | Percentage of Medicare beneficiaries age 65 or older who are chronically disabled, by level and category of disability, 1982-94 | National Long Term Care Survey | Updated with 1999 data and expanded with a change in coding schemes and weights. Data from 1982 are no longer included. | 19A | Age-adjusted percentage of Medicare enrollees age 65 and over who are chronically disabled, by selected characteristics, 1984, 1989, 1994, and 1999 | National Long Term Care Survey | Not updated. | 19A | Age-adjusted percentage of Medicare enrollees age 65 and over who are chronically disabled, by selected characteristics, 1984, 1989, 1994, and 1999 | National Long Term Care Survey | No |
| 18B | Percentage of persons age 70 or older who are unable to perform certain physical functions, by sex, 1984 and 1995 | Supplement on Aging, Second Supplement on Aging | New table. Changed data source from SOA I&II. Different physical functioning measures. | 19B | Percentage of Medicare enrollees age 65 and over who are unable to perform certain physical functions, by sex, 1991 and 2002 | Medicare Current Beneficiary Survey | Updated with 2003 data. The 1991 estimate for women who are unable to stoop/kneel should be changed from 15.0 to 15.1. | 19B | Percentage of Medicare enrollees age 65 and over who are unable to perform certain physical functions, by sex, 1991 and 2003 | Medicare Current Beneficiary Survey | 2004 |

Disability (continued)

| Table # in OA 2000 | Table titles in OA2000 | Data source | Summary of changes between OA2000 and OA2004 | Table # in OA 2004 | Table titles in OA2004 | Data source | Summary of changes between OA2004 and OA2006 | Table # in OA 2006 | Table titles in OA2006 | Data source | Updated since OA2006 |
|--------------------|---|---|--|--------------------|--|-------------------------------------|--|--------------------|--|-------------------------------------|----------------------|
| 18C | Percentage of persons age 70 or older who are unable to perform any one of nine physical functions, by sex and race, 1995 | Supplement on Aging, Second Supplement on Aging | New table. Changed data source from SOA I&II. Different physical functioning measures and race categories. | 19C | Percentage of Medicare enrollees age 65 and over who are unable to perform any one of five physical functions, by selected characteristics, 2002 | Medicare Current Beneficiary Survey | Updated with 2003 data. | 19C | Percentage of Medicare enrollees age 65 and over who are unable to perform any one of five physical functions, by selected characteristics, 2003 | Medicare Current Beneficiary Survey | 2004 |

Sensory Impairments and Oral Health

| | | | | | | | | | | | |
|--|--|--|------|-----|--|----------------------------------|-------------------------|-----|--|----------------------------------|----|
| | | | New. | 16A | Percentage of people age 65 and over who reported having any trouble hearing, any trouble seeing, or no natural teeth, by selected characteristics, 2002 | National Health Interview Survey | Updated with 2004 data. | 16A | Percentage of people age 65 and over who reported having any trouble hearing, any trouble seeing, or no natural teeth, by selected characteristics, 2004 | National Health Interview Survey | No |
| | | | New. | 16B | Percentage of people age 65 and over who reported ever having worn a hearing aid, 2002 | National Health Interview Survey | Updated with 2004 data. | 16B | Percentage of people age 65 and over who reported ever having worn a hearing aid, 2004 | National Health Interview Survey | No |
| | | | New. | 16C | Percentage of people age 65 and over who reported certain conditions among those who reported having trouble seeing, 2002 | National Health Interview Survey | Not updated. | 16C | Percentage of people age 65 and over who reported certain conditions among those who reported having trouble seeing, 2002 | National Health Interview Survey | No |